

## **Statins Linked to Diabetes and Complications in Healthy Adults**

*Author: Troy Brown, RN, Medscape*

Statin use for primary prevention of cardiovascular disease among healthy adults has been linked to an increased risk for diabetes, diabetes complications, and overweight/obesity in a new retrospective cohort study that tracked individuals in a database for an average of 6.5 years.

"Whereas the increased risk of diabetes with statins is well-known, the increased risk of diabetic complications has not been previously described," write the authors, led by Ishak Mansi, MD, from the department of medicine, Veterans Affairs (VA) North Texas Health System, Dallas.

They report their findings in the *Journal of General Internal Medicine* and note that these are among the first data to show a connection between statins and diabetes in a relatively healthy group of people.

"The risk of diabetes with statins has been known, but until now it was thought that this might be due to the fact that people who were prescribed statins had greater medical risks to begin with," said Dr Mansi in a VA statement.

But this current work cannot be used to determine this risk/benefit because of missing information relating to various cardiovascular parameters, he and his colleagues say. Hence further research — including randomized controlled studies for prolonged periods and larger-scale prospective studies — are needed to develop a more complete risk/benefit assessment of statin treatment for primary prevention, they stress.

Asked to comment, Alvin C Powers, MD, director, division of diabetes, endocrinology and metabolism, at Vanderbilt University School of Medicine, Nashville, Tennessee, said: "I think the risk/benefit ratio in people with diabetes and statins remains the same as it was before, and the recommendations per the American Diabetes Association still are relevant."

"[The study] confirmed [an] increased risk for diabetes, and in this case, there were more complications of diabetes in the group taking statins, but it's not clear if that is a result of the statins or just the patient population," he told *Medscape Medical News*.

However, the author of another recent study, which reported the largest risk yet seen for diabetes with statins, Markku Laakso, MD, from the University of Eastern Finland and Kuopio University, has urged caution when considering statin use in primary prevention.

"Statins are not meant to be a treatment for everybody. Especially in women, who are at a lower risk of getting cardiovascular disease, maybe we should be more careful when we start statin treatment?" he said in March, when his work was published.

### **Dose-Response Relationship Observed**

Statin use has long been associated with increased incidence of diabetes, but doctors have always maintained that the benefits of statin use outweigh this risk, particularly in a secondary-prevention population, given the powerful effects of statins in reducing cardiovascular risk.

But data on the long-term effects of these associations in a primary-prevention population are very limited, say Dr Mansi and colleagues. And the relationship between statin use and diabetic complications has not been adequately studied in such individuals, they note.