

Children Today Receive More Than 12 Times as Many Vaccine Doses Than in 1940

By Ethan A. Huff, Natural News

It is a fact that children today are far less able and free to develop their own natural immunity through proper nutrition and normal childhood development. This is due to the fact that the vast majority of them, at least in the industrialized world, are bombarded with an obscene onslaught of vaccines before they are even old enough to attend kindergarten. Data compiled by the *National Vaccine Information Center* (NVIC) reveals that children today receive as many as 49 doses of 14 vaccines before they reach age six, which is roughly 12 times higher than the number of vaccines administered to children back in 1940.

Since 1995, the U.S. *Centers for Disease Control and Prevention* (CDC) has issued yearly revisions to its official childhood immunization schedule, typically adding an increasing amount of vaccines to every new revision. Prior to 1995, CDC issued only periodic updates to its official vaccine schedule, which were typically released every five or six years. But in recent years, the number and frequency of vaccines on the yearly-revised schedule has increased dramatically, as have the number of combination, or multivalent, vaccines being recommended for children.

"Our children are among the most highly vaccinated children in the world, and they are among the most chronically ill and disabled," explains NVIC in a powerful infographic outlining the current vaccine schedule. "Today, the *Centers for Disease Control* admits that 1 child in 6 in America is developmentally delayed. During the past quarter century, the number of children with learning disabilities, ADHD, asthma and diabetes has more than tripled ... (and) the number of doses of vaccines that pediatricians give babies and children under age 6 has more than doubled."

Dramatic rise in brain injuries, neurological damage linked to uptick in vaccines

Interestingly, twice as many children today also suffer from chronic brain and immune system dysfunction compared to the number of children in the 1970s, when roughly half as many vaccines were administered to young children. And the overall percentage of brain-injured children in the 1940s was lower still, as only a few primary vaccines were administered to children at that time. The close connection between the rise in the number of vaccines on the vaccine schedule and the number of brain-injured and neurologically-impaired children, in other words, is difficult to deny, and represents a serious cause for concern.

"Vaccine excipients or ingredients in trace or larger amounts depending on specific vaccine (include) lab altered viruses and bacteria; aluminum; mercury; formaldehyde; phenoxyethanol; gluteraldehyde; sodium borate; sodium chloride; sodium acetate; monosodium glutamate (MSG); hydrochloric acid; hydrogen peroxide; lactose; gelatin; yeast protein; egg albumin; bovine and human serum albumin; antibiotics; (and other) unidentified contaminants," adds the NVIC infographic.

Not only are these detrimental vaccine ingredients a cause for concern, but many of today's vaccines are also given in multiple doses and all at the same time, which was never the case during the 1980s and prior. Children typically received only one, and maybe two, vaccinations during a single visit, and came back on subsequent days to receive the others. Today, however, children sometimes receive up to five vaccinations during a single doctor visit, which exposes them to exponentially higher levels of toxic vaccine components.